

Research

## Purification and partial characterization of bacillocin 490, a novel bacteriocin produced by a thermophilic strain of *Bacillus licheniformis*

Luca Martirani<sup>1</sup>, Mario Varcamonti<sup>1</sup>, Gino Naclerio<sup>2</sup> and Maurilio De Felice\*<sup>1</sup>

Address: <sup>1</sup>Department of General and Environmental Physiology, Section of Microbiology, University Federico II, via Mezzocannone 16, 80134 Naples, Italy and <sup>2</sup>Department of Environmental Sciences and Technologies, University of Molise, via Mazzini 8, 86170 Isernia, Italy

E-mail: Luca Martirani - l.martirani@cecon.it; Mario Varcamonti - varcamon@unina.it; Gino Naclerio - naclerio@unimol.it; Maurilio De Felice\* - defelice@unina.it

\*Corresponding author

Published: 18 April 2002

Received: 1 March 2002

*Microbial Cell Factories* 2002, 1:1

Accepted: 18 April 2002

This article is available from: <http://www.microbialcellfactories.com/content/1/1/1>

© 2002 Martirani et al; licensee BioMed Central Ltd. Verbatim copying and redistribution of this article are permitted in any medium for any purpose, provided this notice is preserved along with the article's original URL.

### Abstract

**Background:** Applications of bacteriocins as food preservatives have been so far limited, principally because of their low antimicrobial activity in foods. Nisin is the only bacteriocin of significant use, but applications are restricted principally because of its very low activity at neutral or alkaline pH. Thus the isolation of new bacteriocins active in foods is desirable.

**Results:** We isolated a *Bacillus licheniformis* thermophilic strain producing a bacteriocin with some novel features, named here bacillocin 490. This bacteriocin was inactivated by pronase E and proteinase K and was active against closely related *Bacillus* spp. both in aerobic and in anaerobic conditions. Bactericidal activity was kept during storage at 4°C and was remarkably stable in a wide pH range. The bacteriocin was partially purified by elution after adhesion to cells of the food-isolated strain *Bacillus smithii* and had a rather low mass (2 KDa). Antimicrobial activity against *B. smithii* was observed also when this organism was grown in water buffalo milk.

**Conclusions:** Bacillocin 490 is a novel candidate as a food anti-microbial agent since it displays its activity in milk, is stable to heat treatment and during storage, is active in a wide pH range and has bactericidal activity also at high temperature. These features may allow the use of bacillocin 490 during processes performed at high temperature and as a complementary antimicrobial agent of nisin against some *Bacillus* spp. in non-acidic foods. The small size suggests its use on solid foods.

### Background

Bacteriocins are ribosomally synthesized antimicrobial peptides produced by a number of different bacteria [1], whose activity is generally directed towards species that are closely related or share the same ecological niche [2,3]. Bacteriocins produced by Gram-positive bacteria have at-

tracted much attention because of their potential use as food preservatives [4]. Many bacteriocins from Gram-positive bacteria, some of which produced by *Bacillus* spp., have been isolated and studied [5–13]. Nisin, a bacteriocin secreted by *Lactococcus lactis*, is commercially used as a food preservative in forty Countries since 1983 [14]. Like

the lactic acid bacteria (LAB), that are the most studied bacteriocin-producing microorganisms, some representatives of *Bacillus* spp., such as *B. subtilis* and *B. licheniformis*, are "generally recognized as safe" (GRAS) bacteria [15].

A wider use of bacteriocins as antimicrobial agents in milk and dairy foods has been hindered so far by the low activity of many known bacteriocins in foods. In this study we report the isolation and partial characterization of a new bacteriocin with promising properties as an antimicrobial agent against *Bacillus* spp. frequently found in non-acidic foods.

## Results

Approximately 23,000 bacterial isolates deriving from dairy foods were tested for their ability to produce antimicrobial activity against Gram-positive and Gram-negative species. We found that 0.48% of tested isolates produce antimicrobial activity against *Salmonella*, while a higher percentage (1.41%) produce anti-*Listeria* activity. The proteinaceous nature of the antimicrobial products was assessed by sensitivity to protease. During the screening, we isolated a strain showing high antimicrobial activity against some adjacent colonies. The producer strain was identified as *Bacillus licheniformis* based on standard phenotypic and metabolic characterization, and 16S ribosomal RNA gene sequence, and named *Bacillus licheniformis* 490/5. The bacteriocin produced, called here bacillocin 490, was completely inactivated by 1 mg of pronase E and proteinase K ml<sup>-1</sup> but not by the same amount of trypsin or chemotrypsin.

*B. licheniformis* 490/5 was shown to be thermophilic with optimal growth rate at 65°C. Its growth was inhibited by a commercial bacitracin species *Staphylococcus epidermidis* but not the resistant one *Escherichia coli*.

The activity range of bacillocin 490 (Table 1) was rather narrow and limited to some Gram-positive bacteria. Highest antimicrobial activity was against *B. stearothermophilus*, *B. smithii*, *B. subtilis* and *B. anthracis*. We also observed some inhibition of *B. cereus* and *B. licheniformis*, very low inhibition of *Listeria innocua* and *S. aureus* and no inhibition of *B. thuringensis* and *Streptococcus thermophilus*. This activity spectrum clearly shows that bacillocin 490 is active principally against species phylogenetically related to the producer strain.

During growth the peak of bacteriocin production occurred in the late logarithmic phase and optimal production was obtained with resting cells at 55°C. Bacteriocin production was increased by 58% and 56% in the presence of 1% (w/v) skimmed milk and gelatin, respectively. This latter result is in agreement with previous observa-

**Table 1: Activity spectrum of bacillocin 490**

Microorganism	Sensitivity*
<i>Pseudomonas aeruginosa</i> ATCC 15442	-
<i>Bacillus licheniformis</i> 5 A2	+
<i>Escherichia coli</i> ATCC 13762	-
<i>Listeria innocua</i> our isolates	+
<i>Staphylococcus aureus</i> ATCC 6538	-
<i>Streptococcus faecalis</i> our isolates	-
<i>Salmonella non tphi</i> our isolates	-
<i>Proteus mirabilis</i> our isolates	-
<i>Staphylococcus epidermidis</i> our isolates	+
<i>Bacillus anthracis</i> 7700	+++
<i>Bacillus subtilis</i> AZ56	++
<i>Bacillus cereus</i> 6A2	+
<i>Bacillus stearothermophilus</i> 9A19	+++
<i>Bacillus smithii</i> PRO/S	+++
<i>Bacillus thuringensis</i>	-
<i>Streptococcus thermophilus</i> ST 11	-


\* Results were identical with plates incubated both in the presence and in the absence of oxygen. See Materials and Methods for other experimental details. (-), not inhibited; (+), (++) , (+++): low, high, very high inhibition, respectively.

tions that, in some cases, solid or semi-solid media (supplemented with gelling agents) increase significantly the amount of bacteriocin production [16–18].

Incubation of *B. smithii* (10<sup>6</sup> bacteria per ml) in the presence of bacillocin 490 (1 µg ml<sup>-1</sup>) at 60°C resulted in 96% killing in 30 minutes, indicating that the bacteriocin has a bactericidal effect.

Bacillocin 490 was purified taking advantage of its ability to bind to the sensitive cells of *B. smithii*. After elution of bound proteins with phosphate buffer, pH 4.5 (see Materials and Methods), a single 2 kDa band showing bactericidal activity was obtained by 10–25% acrylamide SDS-PAGE electrophoresis (data not shown) with a very high yield (60%) and increased specific activity (from 33.7 to 237.8 AU per mg of total proteins) with a purification factor of approximately 7. The latter low value is probably explained by small amount of protein secreted and is in agreement with other bacteriocin purifications [19]. The apparent weight of the bacteriocin was determined by gel-filtration and resulted approximately 2 kDa in agreement with electrophoretic data.

Purified bacillocin 490 was partially characterized. It was equally active both in the presence and in the absence of oxygen, showed great stability in a wide pH range, since it had optimum activity at neutral and moderately basic pH and maintained some 60% of activity after incubation at




**Figure 1**  
Thermal resistance of bacillocin 490 at 100°C. Other experimental details are in the Materials and Methods.

pH 3.5 and 11.0, with 95–100% of activity detected rather uniformly from pH 4.5 to 9.0. This is of great advantage from the industrial point of view, because the pH of many foods varies from very acidic to basic. In this connection it is interesting to notice that nisin, the only bacteriocin so far employed commercially as food additive against Gram-positive bacteria, acts in acidic conditions, as its maximum stability is at pH 2.0, and it is practically not soluble at pH 8.0 [20].

Bacillocin 490 showed high stability at 100°C with 46.4% residual activity after 1 hour of exposure at this temperature (Fig. 1). It also showed remarkable stability at 4°C (no loss of activity after 12-month storage).

Bacillocin 490 was also used in preliminary experiments as an antimicrobial agent against the most sensitive species, *B. smithii*, inoculated in sterile water buffalo milk. Bacillocin 490 was not affected by protein aggregation since no difference of stability was observed when the bacteriocin was dissolved in milk and in buffer. As shown in Fig. 2, we observed 50% growth inhibition of *B. smithii*, after 5-h incubation in milk in the presence of 65 ng of bacillocin 490 ml<sup>-1</sup>. This result indicates that this bacteriocin is suitable for milk-based foods.

Two other bacteriocins found in *B. licheniformis* isolates have been described recently. One, named by the authors Lichenin, displays its activity only in strictly anaerobic conditions and against anaerobic organisms [21]. The other, named BSCY2, has a higher molecular mass with respect to both bacillocin 490 and Lichenin, and is moderately heat stable [22].



**Figure 2**  
Survival of *Bacillus smithii* in milk at 25°C in the absence (open circles) and in the presence (bold circles) of bacillocin 490. See Materials and Methods for experimental details.

## Conclusions

We report here the identification, purification and partial characterization of bacillocin 490, a *B. licheniformis* bacteriocin with very promising features for industrial applications. It displayed bactericidal activity in milk as well as in synthetic media, showed remarkable stability to heat and during storage and was active up to 60°C in a wide pH range. The latter feature may allow the use of bacillocin 490 at high temperature and as a complementary antimicrobial agent of nisin against some *Bacillus* spp. in non-acidic foods. The small size of bacillocin 490 suggests its employment on solid foods.

## Materials and Methods

### Bacterial strains and growth conditions

Bacterial strains used are those listed in Table 1, with the addition of *B. licheniformis* 490/5, the bacteriocin producer. All strains, except *S. thermophilus*, were cultured aerobically in TY medium (16 g tryptone l<sup>-1</sup>, 10 g yeast extract l<sup>-1</sup>, 5 g sodium chloride l<sup>-1</sup>). *S. thermophilus* was grown anaerobically using GasPack apparatus (OXOID, Basingstoke, UK) in HJL broth (3% tryptone, 1% yeast extract, 1% lactose, 0.2% beef extract, 0.5% KH<sub>2</sub>PO<sub>4</sub>, pH 6.5) and plated on solid M17 agar containing 1% lactose.

Solid and soft media were prepared by adding 20 g/liter and 7 g/liter of agar to the broth, respectively.

*Bacillus licheniformis* 490/5 and *Bacillus smithii* PRO/S were isolated from dairy products at 65°C as adjacent colonies on the same plate. Taxonomic classification of both

strains was performed by use of API 50 CHB fermentation strips (bioMérieux, Marcy l'Etoile, France) and by sequence analysis of one of the 16S rRNA genes amplified with two oligonucleotides: P1 (5'-GCGGCGTGCCT AAT-ACATGC) and P2 (5'-CACCTTCCGATACGGCTACC), annealing to nucleotides 40 to 59 and 1532 to 1513, respectively, of *B. subtilis rrmE*.

#### Screening of bacteria producing antimicrobial activities

Milk and dairy products were homogenized, diluted and plated onto various selective and non selective media (PCA, M17, MRS) in different growth conditions (varying temperature and oxygen supply). After incubation, individual clones were replicated in duplicate plates, incubated and, after appearance of colonies, one set of plates was u.v.-254nm irradiated for 15 min to kill the cells and overlaid with 5 ml of 50°C – heated soft agar suspension containing 100 µl of an exponentially growing bacterial culture (10<sup>6</sup> cells per ml) of an indicator strain. Plates were incubated at 37°C for 18 h and inhibition zones were measured. Corresponding producer strains were isolated from the non-irradiated set of plates.

#### Bacteriocin production and detection

*Bacillus licheniformis* 490/5 was grown at 55°C in TY broth to stationary phase, and the culture was centrifuged at 2,500×g for 10 min. 20 µl of cell-free filter-sterilized supernatant were spotted on a sterile susceptibility blank discs (OXOID Basingstoke, UK) placed onto a plate filled with 5 ml of solidified soft agar containing 10<sup>6</sup> cells of *Bacillus smithii* (in 100 µl) used as indicator strain for detection of bacteriocin activity. The plate was incubated at 55°C for 18 h in the presence and in the absence of oxygen (the latter condition was accomplished by incubating the plates in GasPack apparatus) and inhibition areas were measured using a GelDoc 2000 photographic apparatus (BioRad) equipped with MultyAnalyst software (BioRad). One Arbitrary Unit (AU) was defined as the amount of bacteriocin producing 1 mm<sup>2</sup> of inhibition area around the sterile disk.

#### Bacteriocin characterization

To evaluate thermal stability, partially purified bacteriocin was heated at 100°C or exposed at 4°C. Samples were collected and the activity measured at various times and reported as % of values of untreated samples.

Bacteriocin was also treated with pronase E, trypsin, α-chymotrypsin, proteinase K (all purchased from Sigma Chemical Co.). Enzymes were filter-sterilized and added to crude bacteriocin preparations at final concentration of 1 mg/ml. Following incubation at 37°C for 2 h, enzymes were denatured by heating at 100°C for 5 min. Untreated samples incubated in the same conditions were used as

controls. Residual activity of samples was determined as mentioned above.

#### Purification

The bacteriocin was partially purified from 50 ml cultures of *Bacillus licheniformis* 490/5. Cells were grown to stationary phase in TY broth at 55°C and then collected by centrifugation (2,500 × g at 4°C, 10 min). The supernatant was filter-sterilized, ultrafiltered on 1,000 cut-off membrane to concentrate 10-fold and finally mixed with a pellet of pre-washed *B. smithii* cells from 1 liter of an 18 h culture at 55°C. The suspension was gently up-down mixed for 1 h to favour adhesion of the bacteriocin to the cell walls, and then centrifuged 5 min at 700 × g. The supernatant containing unbound proteins was discarded and cells were washed 3 times with 5 ml of 50 mM sodium phosphate buffer, pH 7.0. Finally, bound bacteriocin was eluted with the same buffer at pH 4.5.

#### SDS-polyacrylamide gel electrophoresis

Electrophoresis was performed by standard protocols [23]. For detection of bacteriocin activity after gel separation a previously reported method was used [9].

#### Gel-filtration

To determine its apparent molecular weight, the bacteriocin was loaded on a 1.3 ml G-75 Superdex (Pharmacia) gel filtration column, pre-equilibrated with the elution buffer (50 mM sodium phosphate, pH 7.0). Antimicrobial activity was assayed on each fraction.

#### Stability and antimicrobial activity in milk

For stability studies, equal amounts of bacillocin 490 were added to 10 ml of either 50 mM sodium phosphate buffer pH 7.0 or sterile fresh water buffalo milk and incubated at 25°C. 20 µl of samples were tested for antimicrobial activity each day.

To test the efficacy of bacteriocin against microorganisms present in milk, 10 ml of heat-sterilized fresh water buffalo milk containing 10<sup>6</sup> cells per ml of *Bacillus smithii* strain PRO/S were incubated 5 h at 25°C in the presence of 65 ng of bacteriocin ml<sup>-1</sup>. 100 µl samples were diluted in buffer and plated on TY medium each time to detect cell survival. A sample treated in the same conditions, without bacteriocin, was used as a control. Plates were incubated at 55°C.

#### Acknowledgements

This work belongs to the project "Biotecnologie Mediche ed Agro-alimentari" supported by "MIUR-Ministero dell'Istruzione, dell'Università e della Ricerca" and the "Consorzio Technapoli – Parco Scientifico e Tecnologico dell'Area Metropolitana di Napoli" under responsibility of "CEINGE", Naples, Italy.

The authors wish to thank "Bacillus Genetic Stock Center" that provided some *Bacillus* strains.

## References

1. Tagg JR, Dayani AS, Wannamaker LW: **Bacteriocins of Gram-positive bacteria.** *Bacteriol Rev* 1976, **40**:722-756
2. Klaenhammer TR: **Bacteriocins of Lactic Acid Bacteria.** *Biochimie* 1988, **70**:337-349
3. Klaenhammer TR: **Genetics of bacteriocins produced by Lactic Acid Bacteria.** *FEMS Microbiol Rev* 1993, **12**:39-86
4. Gould G: **Industry perspectives on the use of natural antimicrobial and inhibitors for food applications.** *J Food Prot* 1996, **82**:86
5. Favret ME, Yousten AA: **Thuricin: the bacteriocin produced by *Bacillus thuringensis*.** *J Invertebr Pathol* 1989, **53**:206-216
6. Hyronimus B, Le Marrec C, Urdaci MC: **Coagulin, a bacteriocin-like inhibitory substance produced by *Bacillus coagulans* I<sub>4</sub>.** *J Appl Microbiol* 1998, **85**:42-50
7. Klein C, Entian KD: **Genes involved in self-protection against the lantibiotic subtilin produced by *Bacillus subtilis* ATCC 6633.** *Appl Environ Microbiol* 1994, **60**:2793-2801
8. Le Marrec C, Hyronimus B, Bressollier P, Verneuil B, Urdaci MC: **Biochemical and genetic characterization of Coagulin, a new antilisterial bacteriocin in the pediocin family of bacteriocins, produced by *Bacillus coagulans* I<sub>4</sub>.** *Appl Environ Microbiol* 2000, **66**:5213-5220
9. Naclerio G, Ricca E, Sacco M, De Felice M: **Antimicrobial activity of a newly identified bacteriocin of *Bacillus cereus*.** *Appl Environ Microbiol* 1993, **59**:4313-4316
10. Oscariz JC, Lasa I, Pisabarro AG: **Detection and characterization of cerein 7, a new bacteriocin produced by *Bacillus cereus* with a broad spectrum of activity.** *FEMS Microbiol Letters* 1999, **178**:337-341
11. Piuri M, Sanchez-Rivas C, Ruzal SM: **A novel antimicrobial activity of a *Paenibacillus polymixa* strain isolated from regional fermented sausages.** *Lett Appl Microbiol* 1998, **27**:9-13
12. Sthal S: **A new bacteriocin activity: megacin BII encoded by plasmid PSE 203 in strains of *Bacillus megaterium*.** *Arch Microbiol* 1989, **151**:159-165
13. Zheng G, Slavik MF: **Isolation, partial purification and characterization of a bacteriocin produced by a newly isolated *Bacillus subtilis* strain.** *Lett Appl Microbiol* 1999, **28**:363-367
14. Hoover DG, Steenson LR: **Bacteriocins of Lactic Acid Bacteria.** Academic Press 1993
15. Sharp RJ, Scawen MD, Atkinson T: **Fermentation and downstream processing of *Bacillus*.** In *Biotechnology Handbook: Bacillus* (Edited by: Harwood CR) New York, Plenum Press 1989, 255-292
16. Kelstrup J, Gibbons RS: **Bacteriocins from human and rodent *Streptococci*.** *Arch Oral Biol* 1969, **14**:251-258
17. Setten AM, Vogels GD, De Windt F: **Production and purification of a *Staphylococcus epidermidis* bacteriocin.** *J Bacteriol* 1972, **112**:235-242
18. Piard J-C, Muriana PH, Desmazeaud MJ, Klaenhammer TR: **Purification and partial characterization of Lactacin 481, a lanthionine-containing bacteriocin produced by *Lactococcus lactis* subs. *lactis* CNRZ 481.** *Appl Environ Microbiol* 1992, **58**:279-284
19. Zheng G, Yan LZ, Vederasand JC, Zuber P: **Genes of the SBO-ALB locus of *Bacillus subtilis* are required for production of the antilisterial bacteriocin subtilisin.** *J Bacteriol* 1999, **181**:7346-7355
20. Liu W, Hansen JN: **Some chemical and physical properties of nisin, a small-protein antibiotic produced by *Lactococcus lactis*.** *Appl Environ Microbiol* 1990, **56**:2551-2558
21. Pattanaik P, Kaushik JK, Grover S, Batish VK: **Purification and characterization of a bacteriocin-like compound (Lichenin) produced anaerobically by *Bacillus licheniformis* isolated from water buffalo.** *J Appl Microbiol* 2001, **91**:636-645
22. Chang J-Y, Lee H-H, Kim I-C, Chang H-C: **Characterization of a bacteriocin produced by *Bacillus licheniformis* cy2.** *J Korean Soc. Food Sci. Nutr.* 2001, **30**:410-414
23. Sambrook J, Fritsch EF, Maniatis T: **Molecular Cloning: A Laboratory Manual.** Cold Spring Harbor, Cold Spring Harbor Press 1989

Publish with **BioMed Central** and every scientist can read your work free of charge

"BioMedcentral will be the most significant development for disseminating the results of biomedical research in our lifetime."

Paul Nurse, Director-General, Imperial Cancer Research Fund

Publish with **BMC** and your research papers will be:

- available free of charge to the entire biomedical community
- peer reviewed and published immediately upon acceptance
- cited in PubMed and archived on PubMed Central
- yours - you keep the copyright



Submit your manuscript here:

<http://www.biomedcentral.com/manuscript/>

[editorial@biomedcentral.com](mailto:editorial@biomedcentral.com)