

RESEARCH

Open Access



Lycopene overproduction in *Saccharomyces cerevisiae* through combining pathway engineering with host engineering

Yan Chen^{1,2}, Wenhai Xiao^{1,2*} , Ying Wang^{1,2}, Hong Liu^{1,2}, Xia Li^{1,2} and Yingjin Yuan^{1,2}

Abstract

Background: Microbial production of lycopene, a commercially and medically important compound, has received increasing concern in recent years. *Saccharomyces cerevisiae* is regarded as a safer host for lycopene production than *Escherichia coli*. However, to date, the lycopene yield (mg/g DCW) in *S. cerevisiae* was lower than that in *E. coli* and did not facilitate downstream extraction process, which might be attributed to the incompatibility between host cell and heterologous pathway. Therefore, to achieve lycopene overproduction in *S. cerevisiae*, both host cell and heterologous pathway should be delicately engineered.

Results: In this study, lycopene biosynthesis pathway was constructed by integration of *CrtE*, *CrtB* and *CrtI* in *S. cerevisiae* CEN.PK2. When *YPL062W*, a distant genetic locus, was deleted, little acetate was accumulated and approximately 100 % increase in cytosolic acetyl-CoA pool was achieved relative to that in parental strain. Through screening *CrtE*, *CrtB* and *CrtI* from diverse species, an optimal carotenogenic enzyme combination was obtained, and *CrtI* from *Blakeslea trispora* (*BtCrtI*) was found to have excellent performance on lycopene production as well as lycopene proportion in carotenoid. Then, the expression level of *BtCrtI* was fine-tuned and the effect of cell mating types was also evaluated. Finally, potential distant genetic targets (*YJL064W*, *ROX1*, and *DOS2*) were deleted and a stress-responsive transcription factor *INO2* was also up-regulated. Through the above modifications between host cell and carotenogenic pathway, lycopene yield was increased by approximately 22-fold (from 2.43 to 54.63 mg/g DCW). Eventually, in fed-batch fermentation, lycopene production reached 55.56 mg/g DCW, which is the highest reported yield in yeasts.

Conclusions: *Saccharomyces cerevisiae* was engineered to produce lycopene in this study. Through combining host engineering (distant genetic loci and cell mating types) with pathway engineering (enzyme screening and gene fine-tuning), lycopene yield was stepwise improved by 22-fold as compared to the starting strain. The highest lycopene yield (55.56 mg/g DCW) in yeasts was achieved in 5-L bioreactors. This study provides a good reference of combinatorial engineering of host cell and heterologous pathway for microbial overproduction of pharmaceutical and chemical products.

Keywords: Metabolic engineering, Lycopene, *Saccharomyces cerevisiae*, Heterologous pathway, Synthetic biology

Background

Artificial biosynthetic pathway and host cell are two fundamental elements for microbe-based heterologous biosynthesis of natural products. On one hand, potential metabolic and regulatory issues from host cell play

an important role in pathway productivity [1–4]. One-hundred distant genetic loci that are not directly involved in target pathway were identified to influence carotenoid production significantly in *Saccharomyces cerevisiae* [5]. On the other hand, balanced metabolic flux between modules in target pathway is another important issue to improve pathway performance. Through multivariate-modular optimization of taxadiene metabolic pathway, a 15,000-fold increase in taxadiene titer was observed

*Correspondence: wenhai.xiao@tju.edu.cn

¹ Key Laboratory of Systems Bioengineering (Ministry of Education), Tianjin University, Tianjin 300072, People's Republic of China
Full list of author information is available at the end of the article

in *Escherichia coli* [6]. A “push–pull–block” pathway manipulation strategy significantly enhanced terpenoids production in yeasts [7, 8]. Thus, optimal pathway output can be achieved by means of delicate engineering of both target pathway and host cell [9]. It was reported that bisabolene production in *S. cerevisiae* was increased by 20 times through deleting multiple distant genes related to intracellular mevalonate level and manipulating the expression level of three genes involved in mevalonate (MVA) pathway [10]. Swidah et al. [11] reported that through the combinatorial effects of deletion of *ADHI* to restore redox imbalance, expression of a butanol resistant allele *GCDI*, and manipulation of acetyl-CoA formation module, butanol production in *S. cerevisiae* was increased by more than 30 times. In a word, combinatorial engineering host cell with heterologous pathway offers a promising alternative to achieve better metabolic flux balance and higher output of heterologous pathway.

Lycopene has long been used as functional food, nutraceutical, pharmaceutical and cosmetic due to its antioxidative and anti-cancer activities [12, 13]. Compared to chemical synthesis and extraction from tomatoes, microbial production of lycopene is more economical and sustainable. In recent years, lycopene production was successfully realized in *Blakeslea trispora*, *E. coli* and yeasts. However, regarding to food safety issues, it is controversial to use *B. trispora* or *E. coli* for lycopene synthesis, since *E. coli* would release endotoxin [14] and *B. trispora* requires the addition of cyclase inhibitors [15]. *Saccharomyces cerevisiae* is generally recognized as safe (GRAS), robust and preferred organism for industrial use. To date, lycopene yield in *S. cerevisiae* was increased to 24.41 mg/g DCW with elaborate efforts in directed evolution and copy number variation of *Crt* genes from *Xanthophyllomyces dendrorhous* [16]. However, the lycopene yield was still much lower than that in *E. coli* [17, 18], which did not facilitate downstream extraction process. It was speculated that such low yield might be attributed to the incompatibility between *S. cerevisiae* and the heterologous pathway. Therefore, combinatorial engineering *S. cerevisiae* with a heterologous pathway may offer an effective solution to enhance lycopene yield.

In this study, heterologous carotenogenic pathway and its recruited host *S. cerevisiae* were combinatorially engineered (Fig. 1c). Acetyl-CoA formation was enhanced by the deletion of *YPL062W*. A novel and optimal combination of geranylgeranyl diphosphate synthase (CrtE, or GGPPS), phytoene synthase (CrtB) and phytoene desaturase (CrtI) was generated through related enzyme screening from diverse species. The expression level of *CrtI*, which has significant impact on lycopene production as well as lycopene proportion in carotenoid, was

fine-tuned by varying promoter strength and copy number. The influences of different cell mating types and potential distant genetic loci were also evaluated. Using this combinatorial engineering strategy, we achieved approximately 22-fold improvement (up to 54.63 mg/g DCW) in lycopene yield, which provides a good reference to increase the compatibility between heterologous pathway and host cell for microbial production of valuable molecules.

Methods

Materials

All oligonucleotides were purchased from Invitrogen. Q5 DNA polymerase and *PmeI* restriction endonuclease were purchased from New England Biolabs (MA, USA). DNA purification and plasmids isolation kits were purchased from Tiangen (Beijing, China). CloneJET PCR Cloning Kit was purchased from Fermentas (MD, USA). DNA sequencing was conducted by Genewiz (Beijing, China). Standards of lycopene and phytoene were purchased from Sigma (Sigma-Aldrich, MO, USA). Standards of phytofluene, ζ -carotene and neurosporene were purchased from Express (Beijing, China).

Strains and culture conditions

Escherichia coli DH5 α was used for routine cloning procedures, and was cultivated at 37 °C in Luria–Bertani (LB) medium containing 100 μ g/mL ampicillin for selection. All the yeast strains engineered in this study are based on homologous haploid *S. cerevisiae* strains, CEN.PK2-1C (*MAT α*) or CEN.PK2-1D (*MAT α*). Engineered yeast strains were selected on synthetic complete (SC) medium (0.67 % yeast nitrogen base without amino acids, 2 % glucose, and appropriate amino acid drop-out mix), or YPD medium (2 % peptone, 1 % yeast extract, and 2 % glucose) with 200 μ g/mL geneticin, 300 μ g/mL hygromycin B or 50 μ g/mL bleomycin when needed. YPDG medium, consisting of 2 % glucose (unless otherwise indicated), 2 % peptone, 1 % yeast extract and 1 % D-(+)-galactose, was used for shake-flask fermentations.

For shake-flask fermentation, yeast glycerol-stock was inoculated into a tube containing 2 mL YPD medium for overnight growth, then all the preculture was inoculated into a 250 mL shake-flask containing 25 mL YPD. After growing to the mid-log phase, the seed was transferred to 50 mL fresh YPDG medium at an initial OD₆₀₀ of 0.5 and grown at 30 °C for 48 h.

Construction of plasmids and strains

Saccharomyces cerevisiae strains and plasmids used in this study are summarized in Table 1. All oligonucleotides used for construction of the above plasmids and strains are listed in Additional file 1: Table S1. Construction




Fig. 1 Schematic representation of the engineering strategies for enhanced lycopene production in *S. cerevisiae*. **a** The engineered lycopene biosynthetic pathway in *S. cerevisiae*. Genetic modifications are noted by thick arrows. Red arrows indicate the heterologous lycopene biosynthetic pathway consisting of *CrtE*, *CrtB* and *CrtI*, which starts either from FPP (solid arrow) or directly from IPP and DMAPP (dashed arrow). The enzyme overexpressed in the described lycopene-producing strain is highlighted in blue. **b** Construction of plasmids and integration modules. **c** The rationale of experiment design: combinatorial engineering of host cell and heterologous pathway

procedures of plasmids and integration modules are shown in Fig. 1b. All heterologous genes used for lycopene biosynthesis were codon-optimized and synthesized by Genewiz (Beijing, China) for expression in *S. cerevisiae*. Endogenous truncated 3-hydroxy-3-methylglutaryl coenzyme A reductase (*tHMG1*) [19], promoters, terminators and integration homologous arms (except *TRP1* homologous arm) used in this study were amplified from the genomic DNA of *S. cerevisiae* CEN.PK2-1C. Auxotroph markers (*LEU2*, *HIS3*) and *TRP1* homologous arm were amplified from the genomic DNA of *S. cerevisiae* S288C. Antibiotic markers (*KanMX*, *HphMX* and *BleMX*) were amplified from plasmid pKan, pHph and

pBle (owned by our laboratory), respectively. Overlap extension PCR (OE-PCR) was used to assemble the above parts into modules according to Additional file 1: Figure S1. The resulting modules were purified and cloned into pJET1.2/blunt vector following the protocol of CloneJET PCR Cloning Kit (Fermentas, MD, USA), obtaining plasmid pCY series (Table 1). Finally, all the integration modules were digested from plasmids with *PmeI*, then purified and transformed into yeast for genomic integration using the LiAc/SS carrier DNA/PEG method [20]. For gene deletions, one-step integration of PCR-amplified deletion cassettes was adopted [21]. Gene deletions and genomic integrations were verified by diagnostic PCR.

Table 1 *S. cerevisiae* strains and plasmids used in this study

	Description	Source
Strain name		
CEN.PK2-1C	<i>MATa, ura3-52, trp1-289, leu2-3,112, his3Δ1, MAL2-8C, SUC2</i>	EUROSCARF
CEN.PK2-1D	<i>MATa, ura3-52, trp1-289, leu2-3,112, his3Δ1, MAL2-8C, SUC2</i>	EUROSCARF
SyBE_ Sc14C01	CEN.PK2-1C, <i>Δgal80::HIS3</i>	This study
SyBE_ Sc14C02	CEN.PK2-1C, <i>Δgal1 Δgal7 Δgal10::HIS3</i>	This study
SyBE_ Sc14C06	SyBE_Sc14C01, <i>trp1::TRP1_T_{CYC1}-BtCrtI-P_{GAL10}-P_{GAL1}-PaCrtB-T_{PGK1}, leu2::LEU2_T_{ACT1}-tHMG1-P_{GAL10}-P_{GAL1}-PaCrtE-T_{GPM1}</i>	This study
SyBE_ Sc14C07	SyBE_Sc14C02, <i>trp1::TRP1_T_{CYC1}-BtCrtI-P_{GAL10}-P_{GAL1}-PaCrtB-T_{PGK1}, leu2::LEU2_T_{ACT1}-tHMG1-P_{GAL10}-P_{GAL1}-PaCrtE-T_{GPM1}</i>	This study
SyBE_ Sc14C10	CEN.PK2-1C, <i>Δgal1 Δgal7 Δgal10::HIS3, Δypl062w::KanMX</i>	This study
SyBE_ Sc14C21	SyBE_Sc14C10, <i>trp1::TRP1_T_{CYC1}-AaCrtI-P_{GAL10}-P_{GAL1}-AaCrtB-T_{PGK1}, leu2::LEU2_T_{ACT1}-tHMG1-P_{GAL10}-P_{GAL1}-PaCrtE-T_{GPM1}</i>	This study
SyBE_ Sc14C51	SyBE_Sc14C10, <i>trp1::TRP1_T_{CYC1}-PaCrtI-P_{GAL10}-P_{GAL1}-AaCrtB-T_{PGK1}, leu2::LEU2_T_{ACT1}-tHMG1-P_{GAL10}-P_{GAL1}-PaCrtE-T_{GPM1}</i>	This study
SyBE_ Sc14C52	SyBE_Sc14C10, <i>trp1::TRP1_T_{CYC1}-BtCrtI-P_{GAL10}-P_{GAL1}-AaCrtB-T_{PGK1}, leu2::LEU2_T_{ACT1}-tHMG1-P_{GAL10}-P_{GAL1}-PaCrtE-T_{GPM1}</i>	This study
SyBE_ Sc14C53	SyBE_Sc14C10, <i>trp1::TRP1_T_{CYC1}-AaCrtI-P_{GAL10}-P_{GAL1}-PaCrtB-T_{PGK1}, leu2::LEU2_T_{ACT1}-tHMG1-P_{GAL10}-P_{GAL1}-PaCrtE-T_{GPM1}</i>	This study
SyBE_ Sc14C22	SyBE_Sc14C10, <i>trp1::TRP1_T_{CYC1}-PaCrtI-P_{GAL10}-P_{GAL1}-PaCrtB-T_{PGK1}, leu2::LEU2_T_{ACT1}-tHMG1-P_{GAL10}-P_{GAL1}-PaCrtE-T_{GPM1}</i>	This study
SyBE_ Sc14C23	SyBE_Sc14C10, <i>trp1::TRP1_T_{CYC1}-BtCrtI-P_{GAL10}-P_{GAL1}-PaCrtB-T_{PGK1}, leu2::LEU2_T_{ACT1}-tHMG1-P_{GAL10}-P_{GAL1}-PaCrtE-T_{GPM1}</i>	This study
SyBE_ Sc14C24	SyBE_Sc14C10, <i>trp1::TRP1_T_{CYC1}-AaCrtI-P_{GAL10}-P_{GAL1}-AaCrtB-T_{PGK1}, leu2::LEU2_T_{ACT1}-tHMG1-P_{GAL10}-P_{GAL1}-SaCrtE-T_{GPM1}</i>	This study
SyBE_ Sc14C54	SyBE_Sc14C10, <i>trp1::TRP1_T_{CYC1}-PaCrtI-P_{GAL10}-P_{GAL1}-AaCrtB-T_{PGK1}, leu2::LEU2_T_{ACT1}-tHMG1-P_{GAL10}-P_{GAL1}-SaCrtE-T_{GPM1}</i>	This study
SyBE_ Sc14C55	SyBE_Sc14C10, <i>trp1::TRP1_T_{CYC1}-BtCrtI-P_{GAL10}-P_{GAL1}-AaCrtB-T_{PGK1}, leu2::LEU2_T_{ACT1}-tHMG1-P_{GAL10}-P_{GAL1}-SaCrtE-T_{GPM1}</i>	This study
SyBE_ Sc14C56	SyBE_Sc14C10, <i>trp1::TRP1_T_{CYC1}-AaCrtI-P_{GAL10}-P_{GAL1}-PaCrtB-T_{PGK1}, leu2::LEU2_T_{ACT1}-tHMG1-P_{GAL10}-P_{GAL1}-SaCrtE-T_{GPM1}</i>	This study
SyBE_ Sc14C25	SyBE_Sc14C10, <i>trp1::TRP1_T_{CYC1}-PaCrtI-P_{GAL10}-P_{GAL1}-PaCrtB-T_{PGK1}, leu2::LEU2_T_{ACT1}-tHMG1-P_{GAL10}-P_{GAL1}-SaCrtE-T_{GPM1}</i>	This study
SyBE_ Sc14C26	SyBE_Sc14C10, <i>trp1::TRP1_T_{CYC1}-BtCrtI-P_{GAL10}-P_{GAL1}-PaCrtB-T_{PGK1}, leu2::LEU2_T_{ACT1}-tHMG1-P_{GAL10}-P_{GAL1}-SaCrtE-T_{GPM1}</i>	This study
SyBE_ Sc14C27	SyBE_Sc14C10, <i>trp1::TRP1_T_{CYC1}-AaCrtI-P_{GAL10}-P_{GAL1}-AaCrtB-T_{PGK1}, leu2::LEU2_T_{ACT1}-tHMG1-P_{GAL10}-P_{GAL1}-AfCrtE-T_{GPM1}</i>	This study
SyBE_ Sc14C57	SyBE_Sc14C10, <i>trp1::TRP1_T_{CYC1}-PaCrtI-P_{GAL10}-P_{GAL1}-AaCrtB-T_{PGK1}, leu2::LEU2_T_{ACT1}-tHMG1-P_{GAL10}-P_{GAL1}-AfCrtE-T_{GPM1}</i>	This study
SyBE_ Sc14C58	SyBE_Sc14C10, <i>trp1::TRP1_T_{CYC1}-BtCrtI-P_{GAL10}-P_{GAL1}-AaCrtB-T_{PGK1}, leu2::LEU2_T_{ACT1}-tHMG1-P_{GAL10}-P_{GAL1}-AfCrtE-T_{GPM1}</i>	This study
SyBE_ Sc14C59	SyBE_Sc14C10, <i>trp1::TRP1_T_{CYC1}-AaCrtI-P_{GAL10}-P_{GAL1}-PaCrtB-T_{PGK1}, leu2::LEU2_T_{ACT1}-tHMG1-P_{GAL10}-P_{GAL1}-AfCrtE-T_{GPM1}</i>	This study
SyBE_ Sc14C28	SyBE_Sc14C10, <i>trp1::TRP1_T_{CYC1}-PaCrtI-P_{GAL10}-P_{GAL1}-PaCrtB-T_{PGK1}, leu2::LEU2_T_{ACT1}-tHMG1-P_{GAL10}-P_{GAL1}-AfCrtE-T_{GPM1}</i>	This study
SyBE_ Sc14C29	SyBE_Sc14C10, <i>trp1::TRP1_T_{CYC1}-BtCrtI-P_{GAL10}-P_{GAL1}-PaCrtB-T_{PGK1}, leu2::LEU2_T_{ACT1}-tHMG1-P_{GAL10}-P_{GAL1}-AfCrtE-T_{GPM1}</i>	This study
SyBE_ Sc14C30	SyBE_Sc14C10, <i>trp1::TRP1_T_{CYC1}-AaCrtI-P_{GAL10}-P_{GAL1}-AaCrtB-T_{PGK1}, leu2::LEU2_T_{ACT1}-tHMG1-P_{GAL10}-P_{GAL1}-BtCrtE-T_{GPM1}</i>	This study
SyBE_ Sc14C60	SyBE_Sc14C10, <i>trp1::TRP1_T_{CYC1}-PaCrtI-P_{GAL10}-P_{GAL1}-AaCrtB-T_{PGK1}, leu2::LEU2_T_{ACT1}-tHMG1-P_{GAL10}-P_{GAL1}-BtCrtE-T_{GPM1}</i>	This study
SyBE_ Sc14C61	SyBE_Sc14C10, <i>trp1::TRP1_T_{CYC1}-BtCrtI-P_{GAL10}-P_{GAL1}-AaCrtB-T_{PGK1}, leu2::LEU2_T_{ACT1}-tHMG1-P_{GAL10}-P_{GAL1}-BtCrtE-T_{GPM1}</i>	This study
SyBE_ Sc14C62	SyBE_Sc14C10, <i>trp1::TRP1_T_{CYC1}-AaCrtI-P_{GAL10}-P_{GAL1}-PaCrtB-T_{PGK1}, leu2::LEU2_T_{ACT1}-tHMG1-P_{GAL10}-P_{GAL1}-BtCrtE-T_{GPM1}</i>	This study

Table 1 continued

	Description	Source
SyBE_Sc14C31	SyBE_Sc14C10, <i>trp1::TRP1</i> _T _{CYC1} -PaCrtI-P _{GAL10} -P _{GAL1} -PaCrtB-T _{PGK1} , <i>leu2::LEU2</i> _T _{ACT1} -tHMG1-P _{GAL10} -P _{GAL1} -BtCrtE-T _{GPM1}	This study
SyBE_Sc14C32	SyBE_Sc14C10, <i>trp1::TRP1</i> _T _{CYC1} -BtCrtI-P _{GAL10} -P _{GAL1} -PaCrtB-T _{PGK1} , <i>leu2::LEU2</i> _T _{ACT1} -tHMG1-P _{GAL10} -P _{GAL1} -BtCrtE-T _{GPM1}	This study
SyBE_Sc14C33	SyBE_Sc14C10, <i>trp1::TRP1</i> _T _{CYC1} -AaCrtI-P _{GAL10} -P _{GAL1} -AaCrtB-T _{PGK1} , <i>leu2::LEU2</i> _T _{ACT1} -tHMG1-P _{GAL10} -P _{GAL1} -TmCrtE-T _{GPM1}	This study
SyBE_Sc14C63	SyBE_Sc14C10, <i>trp1::TRP1</i> _T _{CYC1} -PaCrtI-P _{GAL10} -P _{GAL1} -AaCrtB-T _{PGK1} , <i>leu2::LEU2</i> _T _{ACT1} -tHMG1-P _{GAL10} -P _{GAL1} -TmCrtE-T _{GPM1}	This study
SyBE_Sc14C64	SyBE_Sc14C10, <i>trp1::TRP1</i> _T _{CYC1} -BtCrtI-P _{GAL10} -P _{GAL1} -AaCrtB-T _{PGK1} , <i>leu2::LEU2</i> _T _{ACT1} -tHMG1-P _{GAL10} -P _{GAL1} -TmCrtE-T _{GPM1}	This study
SyBE_Sc14C65	SyBE_Sc14C10, <i>trp1::TRP1</i> _T _{CYC1} -AaCrtI-P _{GAL10} -P _{GAL1} -PaCrtB-T _{PGK1} , <i>leu2::LEU2</i> _T _{ACT1} -tHMG1-P _{GAL10} -P _{GAL1} -TmCrtE-T _{GPM1}	This study
SyBE_Sc14C34	SyBE_Sc14C10, <i>trp1::TRP1</i> _T _{CYC1} -PaCrtI-P _{GAL10} -P _{GAL1} -PaCrtB-T _{PGK1} , <i>leu2::LEU2</i> _T _{ACT1} -tHMG1-P _{GAL10} -P _{GAL1} -TmCrtE-T _{GPM1}	This study
SyBE_Sc14C35	SyBE_Sc14C10, <i>trp1::TRP1</i> _T _{CYC1} -BtCrtI-P _{GAL10} -P _{GAL1} -PaCrtB-T _{PGK1} , <i>leu2::LEU2</i> _T _{ACT1} -tHMG1-P _{GAL10} -P _{GAL1} -TmCrtE-T _{GPM1}	This study
SyBE_Sc14C40	SyBE_Sc14C10, <i>leu2::LEU2</i> _T _{CYC1} -RFP	This study
SyBE_Sc14C41	SyBE_Sc14C10, <i>leu2::LEU2</i> _T _{CYC1} -RFP-P _{GAL3}	This study
SyBE_Sc14C42	SyBE_Sc14C10, <i>leu2::LEU2</i> _T _{CYC1} -RFP-P _{GAL7}	This study
SyBE_Sc14C43	SyBE_Sc14C10, <i>leu2::LEU2</i> _T _{CYC1} -RFP-P _{GAL10}	This study
SyBE_Sc14C44	SyBE_Sc14C10, <i>trp1::TRP1</i> _T _{CYC1} -BtCrtI-P _{GAL10} -P _{GAL1} -PaCrtB-T _{PGK1} , <i>leu2::LEU2</i> _T _{CYC1} -BtCrtI-P _{GAL3} -T _{ACT1} -tHMG1-P _{GAL10} -P _{GAL1} -TmCrtE-T _{GPM1}	This study
SyBE_Sc14C45	SyBE_Sc14C10, <i>trp1::TRP1</i> _T _{CYC1} -BtCrtI-P _{GAL10} -P _{GAL1} -PaCrtB-T _{PGK1} , <i>leu2::LEU2</i> _T _{CYC1} -BtCrtI-P _{GAL7} -T _{ACT1} -tHMG1-P _{GAL10} -P _{GAL1} -TmCrtE-T _{GPM1}	This study
SyBE_Sc14C46	SyBE_Sc14C10, <i>trp1::TRP1</i> _T _{CYC1} -BtCrtI-P _{GAL10} -P _{GAL1} -PaCrtB-T _{PGK1} , <i>leu2::LEU2</i> _T _{CYC1} -BtCrtI-P _{GAL7} -T _{ACT1} -tHMG1-P _{GAL10} -P _{GAL1} -TmCrtE-T _{GPM1} , Δ YGLCtau3::HphMX_P _{GAL7} -BtCrtI-T _{CYC1}	This study
SyBE_Sc14D04	CEN.PK2-1D, Δ gal1 Δ gal7 Δ gal10::HIS3, Δ ypl062w::KanMX, <i>trp1::TRP1</i> _T _{CYC1} -BtCrtI-P _{GAL10} -P _{GAL1} -PaCrtB-T _{PGK1}	This study
SyBE_Sc14D05	SyBE_Sc14D04, <i>leu2::LEU2</i> _T _{ACT1} -tHMG1-P _{GAL10} -P _{GAL1} -TmCrtE-T _{GPM1}	This study
SyBE_Sc14D06	SyBE_Sc14D04, <i>leu2::LEU2</i> _T _{CYC1} -BtCrtI-P _{GAL3} -T _{ACT1} -tHMG1-P _{GAL10} -P _{GAL1} -TmCrtE-T _{GPM1}	This study
SyBE_Sc14D07	SyBE_Sc14D04, <i>leu2::LEU2</i> _T _{CYC1} -BtCrtI-P _{GAL7} -T _{ACT1} -tHMG1-P _{GAL10} -P _{GAL1} -TmCrtE-T _{GPM1}	This study
SyBE_Sc14D08	SyBE_Sc14D04, <i>leu2::LEU2</i> _T _{CYC1} -BtCrtI-P _{GAL7} -T _{ACT1} -tHMG1-P _{GAL10} -P _{GAL1} -TmCrtE-T _{GPM1} , Δ YGLCtau3::HphMX_P _{GAL7} -BtCrtI-T _{CYC1}	This study
SyBE_Sc14D10	SyBE_Sc14D04, <i>leu2::LEU2</i> _T _{CYC1} -BtCrtI-P _{GAL7} -T _{ACT1} -tHMG1-P _{GAL10} -P _{GAL1} -TmCrtE-T _{GPM1} , Δ yjl064w::BleMX	This study
SyBE_Sc14D11	SyBE_Sc14D04, <i>leu2::LEU2</i> _T _{CYC1} -BtCrtI-P _{GAL7} -T _{ACT1} -tHMG1-P _{GAL10} -P _{GAL1} -TmCrtE-T _{GPM1} , Δ rox1::BleMX	This study
SyBE_Sc14D12	SyBE_Sc14D04, <i>leu2::LEU2</i> _T _{CYC1} -BtCrtI-P _{GAL7} -T _{ACT1} -tHMG1-P _{GAL10} -P _{GAL1} -TmCrtE-T _{GPM1} , Δ dos2::BleMX	This study
SyBE_Sc14D13	SyBE_Sc14D04, <i>leu2::LEU2</i> _T _{CYC1} -BtCrtI-P _{GAL7} -T _{ACT1} -tHMG1-P _{GAL10} -P _{GAL1} -TmCrtE-T _{GPM1} , Δ rox1::BleMX, Δ dos2::HphMX	This study
SyBE_Sc14D14	SyBE_Sc14D04, <i>leu2::LEU2</i> _T _{CYC1} -BtCrtI-P _{GAL7} -T _{ACT1} -tHMG1-P _{GAL10} -P _{GAL1} -TmCrtE-T _{GPM1} , Δ rox1::BleMX, Δ YPRCdelta15::HphMX_P _{GAL1} -INO2-T _{CPS1}	This study
Plasmid		
pJET1.2/blunt	Blunt-end PCR fragments cloning vector	Fermentas
pCY01	pJET1.2/blunt possessing <i>TRP1</i> homologous arm, T _{CYC1} -AaCrtI-P _{GAL10} -P _{GAL1} -AaCrtB-T _{PGK1}	This study
pCY02	pJET1.2/blunt possessing <i>TRP1</i> homologous arm, T _{CYC1} -PaCrtI-P _{GAL10} -P _{GAL1} -PaCrtB-T _{PGK1}	This study
pCY03	pJET1.2/blunt possessing <i>TRP1</i> homologous arm, T _{CYC1} -BtCrtI-P _{GAL10} -P _{GAL1} -PaCrtB-T _{PGK1}	This study
pCY04	pJET1.2/blunt possessing <i>LEU2</i> homologous arm with <i>LEU2</i> marker, T _{ACT1} -tHMG1-P _{GAL10} -P _{GAL1} -PaCrtE-T _{GPM1}	This study

Table 1 continued

	Description	Source
pCY05	pJET1.2/blunt possessing <i>LEU2</i> homologous arm with <i>LEU2</i> marker, T_{ACT1} - <i>tHMG1</i> - P_{GAL10} - P_{GAL1} - <i>SaCrtE</i> - T_{GPM1}	This study
pCY06	pJET1.2/blunt possessing <i>LEU2</i> homologous arm with <i>LEU2</i> marker, T_{ACT1} - <i>tHMG1</i> - P_{GAL10} - P_{GAL1} - <i>AfCrtE</i> - T_{GPM1}	This study
pCY07	pJET1.2/blunt possessing <i>LEU2</i> homologous arm with <i>LEU2</i> marker, T_{ACT1} - <i>tHMG1</i> - P_{GAL10} - P_{GAL1} - <i>BtCrtE</i> - T_{GPM1}	This study
pCY08	pJET1.2/blunt possessing <i>LEU2</i> homologous arm with <i>LEU2</i> marker, T_{ACT1} - <i>tHMG1</i> - P_{GAL10} - P_{GAL1} - <i>TmCrtE</i> - T_{GPM1}	This study
pCY09	pJET1.2/blunt possessing <i>LEU2</i> homologous arm with <i>LEU2</i> marker, T_{CYC1} - <i>RFP</i>	This study
pCY10	pJET1.2/blunt possessing <i>LEU2</i> homologous arm with <i>LEU2</i> marker, T_{CYC1} - <i>RFP</i> - P_{GAL3}	This study
pCY11	pJET1.2/blunt possessing <i>LEU2</i> homologous arm with <i>LEU2</i> marker, T_{CYC1} - <i>RFP</i> - P_{GAL7}	This study
pCY12	pJET1.2/blunt possessing <i>LEU2</i> homologous arm with <i>LEU2</i> marker, T_{CYC1} - <i>RFP</i> - P_{GAL10}	This study
pCY13	pJET1.2/blunt possessing <i>LEU2</i> homologous arm with <i>LEU2</i> marker, T_{CYC1} - <i>BtCrtI</i> - P_{GAL3} - T_{ACT1} - <i>tHMG1</i> - P_{GAL10} - P_{GAL1} - <i>TmCrtE</i> - T_{GPM1}	This study
pCY14	pJET1.2/blunt possessing <i>LEU2</i> homologous arm with <i>LEU2</i> marker, T_{CYC1} - <i>BtCrtI</i> - P_{GAL7} - T_{ACT1} - <i>tHMG1</i> - P_{GAL10} - P_{GAL1} - <i>TmCrtE</i> - T_{GPM1}	This study
pCY15	pJET1.2/blunt possessing <i>YGLCtau3</i> homologous arm with <i>HphMX</i> marker, P_{GAL7} - <i>BtCrtI</i> - T_{CYC1}	This study
pCY40	pJET1.2/blunt possessing <i>YPRCdelta15</i> homologous arm with <i>HphMX</i> marker, P_{GAL1} - <i>INO2</i> - T_{CPS1}	This study

Assay of extracellular glucose, ethanol, acetate and glycerol

The concentrations of residual glucose, ethanol, acetate and glycerol in the medium were determined by HPLC (Waters Corp., USA) with a refractive index detector. Aminex HPX-87H column (BioRad, CA) was used for separation at the column temperature of 65 °C. 5 mM H₂SO₄ was used as eluent with a flow rate of 0.6 mL/min.

Assay of acetyl-CoA

Cells were sampled during the course of lycopene shake-flask fermentation for acetyl-CoA assay. Acetyl-CoA was extracted as previously described [22] and analyzed by the acetyl-CoenzymeA Assay Kit (Sigma-Aldrich). The acetyl-CoA concentration was normalized by dry cell weight.

Assay of promoter strength

Fluorescence intensity of red fluorescence protein (RFP) was used to characterize the strengths of *GAL3*, *GAL7* and *GAL10* promoters as previously described [23]. The strain without promoter fused with RFP (SyBE_Sc14C40) was used as the negative control. Culturing procedures of all the test strains (SyBE_Sc14C40–SyBE_Sc14C43; Table 1) were the same as lycopene fermentation in shake-flasks. Every 6 h of cultivation, cells were harvested, washed and diluted with phosphate-buffered saline (PBS) into an OD₆₀₀ of 0.3–0.4 for fluorescence assay. RFP fluorescence intensity was detected by SpectraMax M2 microplate reader with excitation and emission wavelengths at 587 and 611 nm, respectively. Promoter strength was determined as the ratio of the fluorescence to OD₆₀₀ for each strain.

Extraction and analysis of carotenoid

Extraction of carotenoid was as described by Xie et al. [24] with some modifications. Briefly, cells harvested from cultures were washed, resuspended in boiling 3 N HCl for 2 min, and cooled in an ice-bath for 3 min. Then, cells debris were washed twice with water, resuspended in acetone containing 1 % BHT (w/v), vortexed with glass beads (425–600 μm, Sigma) until colorless, and followed by centrifugation. The acetone phase containing the extracted carotenoid was filtered for HPLC analysis. A HPLC system (Waters e2695) equipped with a BDS Hypersil C18 column (4.6 × 150 mm, 5 μm) and a UV/VIS detector (Waters 2489) was used to analyze the produced carotenoid. The signals of phytoene, phytofluene, ζ-carotene, neurosporene and lycopene were detected at 287, 349, 401, 440 and 471 nm, respectively [25]. The mobile phase consisted of methanol–acetonitrile–dichloromethane (21:21:8 v/v) with a flow rate of 1 mL/min at 30 °C [26]. Total carotenoid was calculated as the sum of the above carotenoids.

Microscopy

Microscopic analysis was used to investigate lycopene formation of strain SyBE_Sc14C45 during shake-flask fermentation with YPDG medium. SyBE_Sc14C45 cultivated in YPD medium (without galactose) was used as control. After 36 h of cultivation, cells were harvested, washed and diluted with sterile water into an OD₆₀₀ of 5.0. Images were taken with an Olympus CX41 (Olympus, Tokyo, Japan).

Fed-batch fermentation

Strain SyBE_Sc14D14 was selected for fed-batch fermentation. Seed cultures were prepared by inoculating

250 μ L of glycerol-stock into a 250 mL shake-flask containing 25 mL YPD and culturing at 30 °C for 16 h to an OD_{600} of 6–7, and then 15 mL of precultures were inoculated into a 2 L shake-flask containing 400 mL YPD and subcultured for an additional 8 h at 30 °C to an OD_{600} of 5–6. Seed cultures were transferred into a 5 L bioreactor (BaiLun, China) containing 2 L YPD batch medium at a 10 % (v/v) inoculum. Fermentation was carried out at 30 °C with an air flow rate of 1.5 vvm. The dissolved oxygen was kept at 30 % by adjusting the agitation speed from 400 to 700 rpm and pH was controlled at 6.0 by automatic addition of 6 M sodium hydroxide.

According to the employed galactose-inducible system for lycopene biosynthesis, fed-batch fermentation was divided into two stages: cell growth stage and lycopene production stage. During the first stage to achieve maximal cell growth, concentrated glucose solution (500 g/L) was fed periodically into bioreactors to keep the glucose concentration under 2 g/L. In the meanwhile, 100 mL of the concentrated yeast extract solution (400 g/L) was fed periodically into the bioreactor every 10 h. Once cell growth entered stationary phase, glucose and yeast extract feedings were ceased, and 10 g/L of D-(+)-galactose was added to induce lycopene biosynthesis. After the depletion of the residual glucose, cells began to consume ethanol converted by glucose consumption. Ethanol concentration was controlled below 5 g/L by adjusting 100 % ethanol feeding rate until harvest.

Results and discussion

Construction of inducible lycopene biosynthesis pathway

To avoid the potential toxicity of lycopene, genes responsible for carotenoid synthesis were placed under the control of galactose-regulated *GAL* promoters. $\Delta gal1 \Delta gal7 \Delta gal10$ and $\Delta gal80$ were two routine strategies to employ *GAL* promoters, since $\Delta gal1 \Delta gal7 \Delta gal10$ eliminates galactose utilization and $\Delta gal80$ does not require galactose for induction [27]. Here, *S. cerevisiae* SyBE_Sc14C01 ($\Delta gal80$) and SyBE_Sc14C02 ($\Delta gal1 \Delta gal7 \Delta gal10$) were chosen as the host cells for lycopene production. The carotenogenic pathway was constructed by genomic integration of *CrtE*, *CrtB*, *CrtI* and *tHMG1* in the respective hosts (Fig. 1a). As a result, strain SyBE_Sc14C07 ($\Delta gal1 \Delta gal7 \Delta gal10$) with *CrtE* and *CrtB* from *Pantoea agglomerans* and *CrtI* from *Blakeslea trispora* produced 78.8 % higher lycopene yield (4.31 mg/g DCW) than strain SyBE_Sc14C06 ($\Delta gal80$) harboring the same *Crt* genes (2.43 mg/g DCW) after 48 h of shake-flask culture in YPDG medium (Additional file 1: Figure S2). Thus, strain SyBE_Sc14C02 ($\Delta gal1 \Delta gal7 \Delta gal10$) was selected as the host cell for the further study.

Enhancement of acetyl-CoA pool by the deletion of *YPL062W*

$\Delta ypl062w$ (Additional file 1: Table S2) was previously reported to enhance carotenoid production by increasing the intracellular mevalonate level [10], but the mechanism was not clear. In order to testify whether $\Delta ypl062w$ is benefit to lycopene production, *YPL062W* was deleted in SyBE_Sc14C07, generating strain SyBE_Sc14C23. As a result, $\Delta ypl062w$ increased lycopene yield by more than 1.5-fold when cultured in YPDG medium with 2 % glucose (Fig. 2a), which is consistent with previous work [10]. Compared to strain SyBE_Sc14C07 with 0.51 g/L extracellular acetate accumulation, no acetate accumulation was observed in strain SyBE_Sc14C23 (Fig. 2c). Furthermore, SyBE_Sc14C07 and SyBE_Sc14C23 were cultivated in YPDG media with higher (4 %) glucose. As shown in Fig. 2b and d, 10.84 mg/g DCW of lycopene together with very little amount of acetate was detected in SyBE_Sc14C23, whereas only 70 μ g/g DCW of lycopene and up to 5.04 g/L of acetate were obtained in SyBE_Sc14C07. It was also observed that cell growth of strain without $\Delta ypl062w$ was abolished when acetate accumulated up to 1.0 g/L (Fig. 2d), which is in accordance with Thomas et al. [28]. Moreover, when SyBE_Sc14C07 was cultured in YPDG medium with 2 % glucose, different concentrations (0, 0.5, 1.0, 1.5 g/L) of acetic acid were added to the media at the time of 7 h when glucose was exhausted (indicated as arrow in Additional file 1: Figure S3). Cell growth was also abolished when the added acetic acid concentration exceeded 1.0 g/L (Additional file 1: Figure S3A). Lycopene yield was dropped by 98.4 % (from 4.31 mg/g DCW to 67 μ g/g DCW) when additional 0.5 g/L acetic acid was added (Additional file 1: Figure S3B). No lycopene production was detected when 1.0 g/L acetic acid was added (Additional file 1: Figure S3B), suggesting acetate accumulation would be harmful to lycopene biosynthesis. Thus, $\Delta ypl062w$ acted as an important role in *S. cerevisiae* to reduce acetate accumulation.

As is known, acetate is the direct precursor for cytosolic acetyl-CoA. Therefore, we assumed that $\Delta ypl062w$ would enhance cytosolic acetyl-CoA pool from acetate. As expected, the cytosolic acetyl-CoA concentrations in SyBE_Sc14C23 were increased by approximately 100 % than those in SyBE_Sc14C07 at early times (Fig. 2e, f). When lycopene was rapidly accumulated, SyBE_Sc14C23 demonstrated the same cytosolic acetyl-CoA level but higher lycopene production as compared to SyBE_Sc14C07 (Fig. 2a, b, e, f). This result suggested the expansion on lycopene production might be derived from the increase in acetyl-CoA supplement, since the intracellular mevalonate level of $\Delta ypl062w$ strain was significantly increased during terpenoid production [10]. Therefore,




Fig. 2 The effect of $\Delta ypl062w$ on lycopene production. *S. cerevisiae* SyBE_Sc14C07 and SyBE_Sc14C23 were cultivated in YPDG media containing different concentrations of glucose (2 %, left side; 4 %, right side), respectively, in shake-flasks for analysis of lycopene production (a, b), acetate accumulation (c, d) and cytosolic acetyl-CoA level (e, f). The error bars represent standard deviation calculated from triplicate experiments

$\Delta ypl062w$ enhanced “trapping” the carbon from acetate accumulation toward acetyl-CoA, which consequently improved MVA pathway flux and further lycopene production. Our findings made us a better understanding of the effect of $\Delta ypl062w$ on MVA pathway.

Optimal combination of CrtE, CrtB and CrtI by screening enzymes from diverse sources

It is known that enzymes from different organisms often vary in catalytic activities [29, 30]. Besides, host cell compatibility may also affect the optimal performance

of heterologous enzymes [16, 31]. Thus, screening enzymes from diverse sources offers an effective strategy to increase the productivity of heterologous pathway in specific host [29, 32]. To date, most of the carotenogenic genes employed in heterologous biosynthesis were derived from *Pantoea*, *Paracoccus* or *Xanthophyllomyces* species [33, 34]. However, in *S. cerevisiae*, the currently reported *Crt* genes for high-level carotenoid production were only from *X. dendrorhous* [16, 24, 35]. In this study, we aimed to rebuild a carotenogenic pathway with high productivity in *S. cerevisiae* by screening enzymes

(CrtE, CrtB, and CrtI) from some other species except *X. dendrorhous*. Five CrtEs originated from *P. agglomerans* (PaCrtE), *Sulfolobus acidocaldarius* (SaCrtE), *Archaeoglobus fulgidus* (AfCrtE), *B. trispora* (BtCrtE) and *Taxus x media* (TmCrtE), two CrtBs from *P. agglomerans* (PaCrtB) and *Paracoccus* sp. (formerly *Agrobacterium aurantiacum*) (AaCrtB), and three CrtIs from *P. agglomerans* (PaCrtI), *Paracoccus* sp. (AaCrtI) and *B. trispora* (BtCrtI) were selected for carotenoid biosynthesis. As illustrated in Fig. 3, thirty lycopene-producing strains were constructed and their production was investigated. Consequently, the lycopene yield in strain SyBE_Sc14C35 harboring the best enzyme combination (TmCrtE, PaCrtB and BtCrtI) was increased by 7.5-fold, up to 36.75 mg/g DCW, and the lycopene proportion in carotenoid was 64.11 % (Fig. 3). This strain was used as the candidate for the further optimization.

In general, CrtE, one of the rate-limiting enzymes in carotenoid pathway [36], was found to be crucial to the production yield of overall carotenoid. As shown in Fig. 3, strains harboring AfCrtE, BtCrtE or TmCrtE showed much higher yield of total carotenoid than that harboring PaCrtE or SaCrtE. TmCrtE was reported to possess a larger pocket and higher affinity for farnesyl diphosphate (FPP) binding than other GGPPs [30], thus it was reasonable that strain with TmCrtE achieved high carotenoid yield. CrtE from *B. trispora* [37] was firstly well expressed in *S. cerevisiae* and achieved quite high yield of overall carotenoid, demonstrating that BtCrtE would be a promising GGPPS candidate for further study. AfCrtE can directly utilize dimethylallyl pyrophosphate (DMAPP)/isopentenyl pyrophosphate (IPP) to synthesize geranylgeranyl diphosphate (GGPP) and thus

avoid competing FPP for sterols biosynthesis [38], which might explain the improved carotenoid production catalyzed by AfCrtE. In addition to AfCrtE, SaCrtE was also a bifunctional FPP/GGPP synthase [39] and has been demonstrated to increase diterpenoids production [40, 41]. But it was difficult to interpret the low carotenoid yield by SaCrtE according to our current data, which might be ascribed to the insufficient expression of SaCrtE or its incompatibility with CrtB and CrtI.

Phytoene, synthesized by CrtB from GGPP, is the first intermediate of the carotenoid pathway. Then lycopene was generated by CrtI through four successive dehydrogenation steps from phytoene (Fig. 1a). As illustrated in Fig. 3, rather than lycopene, phytoene was one of the major components of the total carotenoid in most of our engineered strains, indicating that CrtI-catalyzed conversion from phytoene to lycopene was another rate-limiting step, which is consistent with previous reports [16, 35]. To be noted, all the strains harboring BtCrtI showed much better performance than strains harboring PaCrtI or AaCrtI, irrespective of lycopene yield or proportion in carotenoid (Fig. 3). BtCrtI, an enzyme from eukaryotic organisms, was firstly well expressed in *S. cerevisiae* and found to be more suitable for high-level conversion from phytoene to lycopene in *S. cerevisiae* according to our results. The molecular mechanism for its high efficiency in conversion from phytoene to lycopene is an interesting topic in the field in future.

Fine-tuning of BtCrtI and selection of homologous haploid yeast hosts

Despite the improved lycopene yield in SyBE_Sc14C35, approximately 26 % of the total carotenoid was phytoene, which indicated that the phytoene conversion




Fig. 3 Combinatorial optimization of CrtE, CrtB and CrtI from diverse species. Thirty lycopene-producing strains were constructed by screening enzymes from various sources and tested for lycopene production. Pa, *Pantoea agglomerans*; Sa, *Sulfolobus acidocaldarius*; Af, *Archaeoglobus fulgidus*; Bt, *Blakeslea trispora*; Tm, *Taxus x media*; Aa, *Paracoccus* sp. (formerly *Agrobacterium aurantiacum*). The error bars represent standard deviation calculated from triplicate experiments

directed by *BtCrtI* was not efficient enough. In order to achieve higher lycopene proportion, the expression level of *BtCrtI* needs to be fine-tuned. Here, *BtCrtI* was fine-tuned by adjusting different promoters and integration copy numbers. The strengths of *GAL* promoters used in this step were characterized in strain SyBE_Sc14C10 (CEN.PK2-1C, $\Delta gal1 \Delta gal7 \Delta gal10, \Delta yjl062w$), and P_{GAL3} was found to be the weakest one (Additional file 1: Figure S4). When one additional copy of *BtCrtI* under the control of P_{GAL3} or P_{GAL7} was integrated in strain SyBE_Sc14C35, the lycopene proportion in the resulting strain SyBE_Sc14C44 or SyBE_Sc14C45 was increased by 17.9 % (from 64.11 to 75.58 %) or 35.2 % (up to 86.68 %), respectively (Fig. 4). Moreover, a slight decrease in lycopene yield was observed after another additional copy of P_{GAL7} -*BtCrtI* integrated in SyBE_Sc14C45 (Fig. 4), which is similar to recent report [16]. To be noted, although the lycopene yield was modestly increased, the total carotenoid yield was decreased obviously after fine-tuning steps for higher lycopene proportion (Fig. 4). As shown in Additional file 1: Figure S5, most of the lycopene was accumulated in cell membrane, which is consistent with early reports [42, 43]. Rapid lycopene accumulation in cell membrane would lead to membrane stress or cell toxicity, which might explain the significant decrement in the content of total carotenoid after fine-tuning. Therefore, increasing lycopene tolerance in *S. cerevisiae* would be an effective direction to improve lycopene yield.

Haploid cell type was reported to have significant influence on heterologous terpenoid production [44]. Jackson et al. [45] found that *MATa* strain was more suitable to

produce epi-cedrol than *MAT α* strain, since *MATa* strain might synthesize more FPP for prenylation of the mating pheromone [46]. However, cell mating types did not have significant effect on linalool production [47]. Here, lycopene production in CEN.PK2-1C (*MATa*) and CEN.PK2-1D (*MAT α*) with the same genetic modifications was evaluated, respectively. The strains (SyBE_Sc14D05–SyBE_Sc14D08) derived from CEN.PK2-1D (*MAT α*) achieved 12–15 % higher lycopene yield than strains (SyBE_Sc14C35, SyBE_Sc14C44–SyBE_Sc14C46) from CEN.PK2-1C (*MATa*) (Fig. 4), suggesting that *MAT α* strain was preferred in the case of lycopene production. Finally, the strain SyBE_Sc14D07 (*MAT α*) achieved lycopene yield of 46.26 mg/g DCW with a proportion of 82.36 % (Fig. 4).

Effects of distant genetic loci on lycopene production

As the engineered metabolic pathways were highly inter-connected with the rest of cellular metabolism and tightly regulated [48], distantly located genetic loci in host cell could also have potential interactions with target pathway. As $\Delta rox1, \Delta dos2$ and $\Delta yjl064w$ (Additional file 1: Table S2) were proved to greatly benefit carotenoid production [5, 10], these three distant genetic loci were knocked out individually in strain SyBE_Sc14D07. As shown in Fig. 5, both $\Delta rox1$ and $\Delta dos2$ conferred a modest increase (8.7 and 5.7 %, respectively) in lycopene production as expected, while $\Delta yjl064w$ led to 18.2 % decreased lycopene yield as compared to SyBE_Sc14D07. Moreover, the combination of $\Delta rox1$ and $\Delta dos2$ did not show a synergistic effect on lycopene production (Fig. 5), which is inconsistent with the results obtained by Triikka




Fig. 4 Carotenoid production by fine-tuning of *BtCrtI* and selecting homologous haploid strains. *BtCrtI* expression was fine-tuned by adjusting copy number and promoter strength, and haploid cells with different mating types (*a, a*) were compared as well for carotenoid production. The error bars represent standard deviation calculated from triplicate experiments




Fig. 5 The effects of distant genetic loci on lycopene production. Three gene-deletion targets (*ROX1, DOS2*, and *YJL064W*) and one overexpression target (*INO2*) were investigated in *S. cerevisiae* SyBE_Sc14D07 for lycopene production. The error bars represent standard deviation calculated from triplicate experiments

et al. [5]. These inconsistencies might be attributed to that the impact of perturbations in one strain may not be directly applied to another strain with a modified genetic background [1]. Consequently, a relatively higher lycopene yield of 50.28 mg/g DCW was obtained in SyBE_Sc14D11 with $\Delta rox1$.

As shown in Additional file 1: Figure S5, lycopene tended to accumulate in cell membrane and thus cause cell toxicity. Here, *INO2* (Additional file 1: Table S2), an endogenous transcription factor related to cellular stress response [49, 50], was overexpressed in SyBE_Sc14D11. As a result, a lycopene yield of 54.63 mg/g DCW, the highest yield reported, was obtained in the resulting strain SyBE_Sc14D14 in shake-flask cultivation (Fig. 5). *INO2* was reported to alleviate alkanes (C9–C11) toxicity by regulating genes associated with efflux pumps, stress response, lipid metabolism and ergosterol biosynthesis in *S. cerevisiae* [51]. High-level expression of *INO2* was proved to up-regulate phospholipid and sterol biosynthesis [52]. Therefore, increasing lycopene tolerance through modifying membrane components (i.e. lipid, and ergosterol) may be the main reason for the improvement of lycopene yield via *INO2*. Therefore, *INO2* was identified as a novel target for lycopene production in *S. cerevisiae*. Strain SyBE_Sc14D14 was chosen for fed-batch fermentation.

Lycopene overproduction in fed-batch fermentation

To evaluate the production performance of the engineered strain SyBE_Sc14D14, fed-batch fermentation was performed at a 2 L scale using YPD as the medium (Fig. 6). Based on carbon restriction strategy, trace amount of acetate was observed during the whole process (Additional file 1: Figure S6). Eventually, a total carotenoid titer of 1.81 g/L (60.94 mg/g DCW), consisting of 3.99 % of phytoene, 4.87 % of neurosporene and 91.14 % of lycopene, was obtained after 120 h of cultivation (Fig. 6). Lycopene yield of 55.56 mg/g DCW achieved in our work was the highest yield in yeast strains to date. However, the lycopene titer (1.65 g/L) was just similar to previous work by Xie et al. [16]. This is due to the relative low cell density, since the highest OD_{600} only reached 106 throughout the fermentation. Additionally, excessive accumulations of ethanol and glycerol were also observed during cell growth phase (Fig. 6a; Additional file 1: Figure S6), which competed carbon flow from biomass synthesis and implied redox imbalance of our engineered strain [53]. In future, to limit glucose below 0.5 g/L in growth phase will greatly increase cell density as well as reduce by-product (ethanol or glycerol) accumulation. Moreover, intracellular and extracellular metabolomics analysis will be an efficient way to find some biomarkers for batch media or feeding solution optimization. Off-gas analysis will also be promising for more precise process control. As recent




Fig. 6 Lycopene production in fed-batch fermentation. **a** Profile of glucose, ethanol, cell density and lycopene accumulation of strain SyBE_Sc14D14 during fed-batch fermentation. **b** Percentage ratios of the produced carotenoid composition at 120 h. The red liquid in the bottles was the lycopene fermentation broth. The error bars represent standard deviation calculated from duplicate experiments

studies in media optimization have demonstrated great potential in lycopene overproduction [16, 54], we believe that lycopene production by our engineered strain would be further improved by continuous efforts in both metabolic engineering and fermentation optimization.

Conclusions

In this work, lycopene overproduction was realized by combinatorial engineering of *S. cerevisiae* and lycopene biosynthesis pathway. Extracellular acetate accumulation was reduced and cytosolic acetyl-CoA pool was enhanced through the deletion of *YPL062W*. A novel and optimal combination of *CrtE*, *CrtB* and *CrtI* was obtained by screening enzymes from diverse sources. It was also found that *CrtI* from *B. trispora* had significant influence on lycopene yield as well as proportion in carotenoid. The proportion of lycopene was significantly increased via fine-tuning of *CrtI*. Then the effects of cell mating types, several potential distant targets (*YJL064 W*, *ROX1*, and *DOS2*), and *INO2*, a stress-related transcription factor, were also investigated. Lycopene yield was stepwise improved by approximately 22-fold as compared to the starting strain. The highest reported lycopene yield (55.56 mg/g DCW) and titer (1.65 g/L) were achieved in 5-L bioreactors, providing a good example for microbial overproduction of pharmaceutical and chemical products through combinatorial engineering of host cell and heterologous pathway.

Additional file

Additional file 1. This file consists of two supplemental tables and seven supplemental figures. **Table S1.** Oligonucleotides used in this study.

Table S2. Distant genetic loci involved in this study. **Figure S1.** Integration modules constructed in this study. **Figure S2.** Comparison of *Δgal1 Δgal7 Δgal10* strain and *Δgal80* strain on lycopene production. **Figure S3.** The effects of acetic acid addition on cell growth (A) and lycopene production (B) in strain SyBE_Sc14C07. **Figure S4.** Time course of promoter strengths. **Figure S5.** Microscopic images of lycopene-producing strain. **Figure S6.** Profile of glycerol and acetate accumulation. **Figure S7.** Sequences of codon-optimized genes.

Abbreviations

MVA: mevalonate; HMG-CoA: 3-hydroxy-3-methylglutaryl coenzyme A; IPP: isopentenyl pyrophosphate; DMAPP: dimethylallyl pyrophosphate; FPP: farnesyl diphosphate; GGPP: geranylgeranyl diphosphate; tHMG1: truncated 3-hydroxy-3-methylglutaryl coenzyme A reductase; CrE: geranylgeranyl diphosphate synthase (GGPPS); CrTb: phytoene synthase; CrTl: phytoene desaturase; Aa: *Paracoccus* sp. (formerly *Agrobacterium aurantiacum*); Pa: *Pantoea agglomerans*; Sa: *Sulfolobus acidocaldarius*; Af: *Archaeoglobus fulgidus*; Bt: *Blakeslea trispora*; Tm: *Taxus x media*.

Authors' contributions

YC, WX and YY conceived of the study and drafted the manuscript. YC carried out the molecular genetic studies. YC and WX carried out the fed-batch fermentation experiments. YW participated in design and coordination of the study and helped to draft the manuscript. HL and XL participated in HPLC analysis and data processing. YY supervised the whole research and revised the manuscript. All authors read and approved the final manuscript.

Author details

¹ Key Laboratory of Systems Bioengineering (Ministry of Education), Tianjin University, Tianjin 300072, People's Republic of China. ² SynBio Research Platform, Collaborative Innovation Center of Chemical Science and Engineering (Tianjin), School of Chemical Engineering and Technology, Tianjin University, Tianjin 300072, People's Republic of China.

Acknowledgements

The authors are grateful for the financial support from the Ministry of Science and Technology of China ("863" Program: 2012AA02A701) and the International S&T Cooperation Program of China (2015DFA00960).

Competing interests

The authors declare that they have no competing interests.

Availability of data and material

The material and data supporting their findings can be found in the main paper and the Additional file 1.

Funding

The Ministry of Science and Technology of China ("863" Program: 2012AA02A701) and the International S&T Cooperation Program of China (2015DFA00960) supported this work.

Received: 28 February 2016 Accepted: 6 June 2016

Published online: 21 June 2016

References

- Alper H, Jin YS, Moxley JF, Stephanopoulos G. Identifying gene targets for the metabolic engineering of lycopene biosynthesis in *Escherichia coli*. *Metab Eng*. 2005;7(3):155–64.
- Alper H, Miyaoku K, Stephanopoulos G. Construction of lycopene-overproducing *E. coli* strains by combining systematic and combinatorial gene knockout targets. *Nat Biotechnol*. 2005;23(5):612–6.
- Brochado AR, Patil KR. Model-guided identification of gene deletion targets for metabolic engineering in *Saccharomyces cerevisiae*. *Methods Mol Biol*. 2014;1152:281–94.
- Jin YS, Stephanopoulos G. Multi-dimensional gene target search for improving lycopene biosynthesis in *Escherichia coli*. *Metab Eng*. 2007;9(4):337–47.
- Trikka FA, Nikolaidis A, Athanasakoglou A, Andreadelli A, Ignea C, Kotta K, Argiriou A, Kampranis SC, Makris AM. Iterative carotenogenic screens identify combinations of yeast gene deletions that enhance sciaene production. *Microb Cell Fact*. 2015;14:60.
- Ajikumar PK, Xiao WH, Tyo KE, Wang Y, Simeon F, Leonard E, Mucha O, Phon TH, Pfeifer B, Stephanopoulos G. Isoprenoid pathway optimization for Taxol precursor overproduction in *Escherichia coli*. *Science*. 2010;330(6000):70–4.
- Chen Y, Daviet L, Schalk M, Siewers V, Nielsen J. Establishing a platform cell factory through engineering of yeast acetyl-CoA metabolism. *Metab Eng*. 2013;15:48–54.
- Lv X, Xie W, Lu W, Guo F, Gu J, Yu H, Ye L. Enhanced isoprene biosynthesis in *Saccharomyces cerevisiae* by engineering of the native acetyl-CoA and mevalonic acid pathways with a push-pull-restrain strategy. *J Biotechnol*. 2014;186:128–36.
- Ma T, Deng Z, Liu T. Microbial production strategies and applications of lycopene and other terpenoids. *World J Microbiol Biotechnol*. 2016;32(1):15.
- Özaydin B, Burd H, Lee TS, Keasling JD. Carotenoid-based phenotypic screen of the yeast deletion collection reveals new genes with roles in isoprenoid production. *Metab Eng*. 2013;15:174–83.
- Swidah R, Wang H, Reid PJ, Ahmed HZ, Pisanelli AM, Persaud KC, Grant CM, Ashe MP. Butanol production in *S. cerevisiae* via a synthetic ABE pathway is enhanced by specific metabolic engineering and butanol resistance. *Biotechnol Biofuels*. 2015;8:97.
- Sies H, Stahl W. Lycopene: antioxidant and biological effects and its bio-availability in the human. *Proc Soc Exp Biol Med*. 1998;218:121–4.
- Giovannucci E, Ascherio A, Rimm EB, Stampfer MJ, Colditz GA, Willett WC. Intake of carotenoids and retinol in relation to risk of prostate cancer. *J Natl Cancer Inst*. 1995;87(23):1767–76.
- Ray BL, Raetz CR. The biosynthesis of gram-negative endotoxin. A novel kinase in *Escherichia coli* membranes that incorporates the 4'-phosphate of lipid A. *J Biol Chem*. 1987;262(3):1122–8.
- Mantzouridou F, Tsimidou MZ. Lycopene formation in *Blakeslea trispora*. Chemical aspects of a bioprocess. *Trends Food Sci Technol*. 2008;19(7):363–71.
- Xie W, Lv X, Ye L, Zhou P, Yu H. Construction of lycopene-overproducing *Saccharomyces cerevisiae* by combining directed evolution and metabolic engineering. *Metab Eng*. 2015;30:69–78.
- Kim YS, Lee JH, Kim NH, Yeom SJ, Kim SW, Oh DK. Increase of lycopene production by supplementing auxiliary carbon sources in metabolically engineered *Escherichia coli*. *Appl Microbiol Biotechnol*. 2011;90(2):489–97.
- Zhu F, Lu L, Fu S, Zhong X, Hu M, Deng Z, Liu T. Targeted engineering and scale up of lycopene overproduction in *Escherichia coli*. *Process Biochem*. 2015;50(3):341–6.
- Donald KA, Hampton RY, Fritz IB. Effects of overproduction of the catalytic domain of 3-hydroxy-3-methylglutaryl coenzyme A reductase on squalene synthesis in *Saccharomyces cerevisiae*. *Appl Environ Microbiol*. 1997;63:3341–4.
- Gietz RD, Schiestl RH. High-efficiency yeast transformation using the LiAc/SS carrier DNA/PEG method. *Nat Protoc*. 2007;2(1):31–4.
- Longtine MS, McKenzie IIIA, Demarini DJ, Shah NG, Wach A, Brachat A, Philippsen P, Pringle JR. Additional modules for versatile and economical PCR-based gene deletion and modification in *Saccharomyces cerevisiae*. *Yeast*. 1998;14(10):953–61.
- Lian J, Si T, Nair NU, Zhao H. Design and construction of acetyl-CoA over-producing *Saccharomyces cerevisiae* strains. *Metab Eng*. 2014;24:139–49.
- Cao YX, Xiao WH, Liu D, Zhang JL, Ding MZ, Yuan YJ. Biosynthesis of odd-chain fatty alcohols in *Escherichia coli*. *Metab Eng*. 2015;29:113–23.
- Xie W, Liu M, Lv X, Lu W, Gu J, Yu H. Construction of a controllable beta-carotene biosynthetic pathway by decentralized assembly strategy in *Saccharomyces cerevisiae*. *Biotechnol Bioeng*. 2014;111(1):125–33.
- Britton G. General carotenoid methods. *Methods Enzymol*. 1985;111(4):113–49.

26. Zhao J, Li Q, Sun T, Zhu X, Xu H, Tang J, Zhang X, Ma Y. Engineering central metabolic modules of *Escherichia coli* for improving beta-carotene production. *Metab Eng*. 2013;17:42–50.
27. Westfall PJ, Pitera DJ, Lenihan JR, Eng D, Woollard FX, Regentin R, Horning T, Tsuruta H, Melis DJ, Owens A, et al. Production of amorphaadiene in yeast, and its conversion to dihydroartemisinic acid, precursor to the anti-malarial agent artemisinin. *Proc Natl Acad Sci USA*. 2012;109(3):E111–8.
28. Thomas KC, Hynes SH, Ingledew WM. Influence of medium buffering capacity on inhibition of *Saccharomyces cerevisiae* growth by acetic and lactic acids. *Appl Environ Microbiol*. 2002;68(4):1616–23.
29. Chang JJ, Thia C, Lin HY, Liu HL, Ho FJ, Wu JT, Shih MC, Li WH, Huang CC. Integrating an algal β -carotene hydroxylase gene into a designed carotenoid-biosynthesis pathway increases carotenoid production in yeast. *Bioresour Technol*. 2015;184:2–8.
30. Ding MZ, Yan HF, Li LF, Zhai F, Shang LQ, Yin Z, Yuan YJ. Biosynthesis of Taxadiene in *Saccharomyces cerevisiae*: selection of geranylgeranyl diphosphate synthase directed by a computer-aided docking strategy. *PLoS ONE*. 2014;9(10):e109348.
31. Greene JJ. Host cell compatibility in protein expression. *Methods Mol Biol*. 2004;267:3–14.
32. Sarria S, Wong B, Garcia Martin H, Keasling JD, Peralta-Yahya P. Microbial synthesis of pinene. *ACS Synth Biol*. 2014;3(7):466–75.
33. Misawa N, Shimada H. Metabolic engineering for the production of carotenoids in non-carotenogenic bacteria and yeasts. *J Biotechnol*. 1997;59(3):169–81.
34. Mata-Gomez LC, Montanez JC, Mendez-Zavala A, Aguilar CN. Biotechnological production of carotenoids by yeasts: an overview. *Microb Cell Fact*. 2014;13:12.
35. Verwaal R, Wang J, Meijnen JP, Visser H, Sandmann G, van den Berg JA, van Ooyen AJ. High-level production of beta-carotene in *Saccharomyces cerevisiae* by successive transformation with carotenogenic genes from *Xanthophyllomyces dendrorhous*. *Appl Environ Microbiol*. 2007;73(13):4342–50.
36. Yoon SH, Kim JE, Lee SH, Park HM, Choi MS, Kim JY, Lee SH, Shin YC, Keasling JD, Kim SW. Engineering the lycopene synthetic pathway in *E. coli* by comparison of the carotenoid genes of *Pantoea agglomerans* and *Pantoea ananatis*. *Appl Microbiol Biotechnol*. 2007;74(1):131–9.
37. Sun J, Sun XX, Tang PW, Yuan QP. Molecular cloning and functional expression of two key carotene synthetic genes derived from *Blakeslea trispora* into *E. coli* for increased β -carotene production. *Biotechnol Lett*. 2012;34(11):2077–82.
38. Wang CW, Oh MK, Liao JC. Engineered isoprenoid pathway enhances astaxanthin production in *Escherichia coli*. *Biotechnol Bioeng*. 1999;62(2):235–41.
39. Ohnuma SI, Suzuki M, Nishino T. Archaeobacterial ether-linked lipid biosynthetic gene. Expression cloning, sequencing, and characterization of geranylgeranyl-diphosphate synthase. *J Biol Chem*. 1994;269(20):14792–7.
40. Dai Z, Liu Y, Huang L, Zhang X. Production of miltiradiene by metabolically engineered *Saccharomyces cerevisiae*. *Biotechnol Bioeng*. 2012;109(11):2845–53.
41. Engels B, Dahm P, Jennewein S. Metabolic engineering of taxadiene biosynthesis in yeast as a first step towards Taxol (Paclitaxel) production. *Metab Eng*. 2008;10(3–4):201–6.
42. Gruszecki WI, Strzalka K. Carotenoids as modulators of lipid membrane physical properties. *Biochim Biophys Acta*. 2005;1740(2):108–15.
43. Verwaal R, Jiang Y, Wang J, Daran JM, Sandmann G, van den Berg JA, van Ooyen AJ. Heterologous carotenoid production in *Saccharomyces cerevisiae* induces the pleiotropic drug resistance stress response. *Yeast*. 2010;27(12):983–98.
44. Ohto C, Muramatsu M, Obata S, Sakuradani E, Shimizu S. Overexpression of the gene encoding HMG-CoA reductase in *Saccharomyces cerevisiae* for production of prenol alcohols. *Appl Microbiol Biotechnol*. 2009;82(5):837–45.
45. Jackson BE, Hart-Wells EA, Matsuda SP. Metabolic engineering to produce sesquiterpenes in yeast. *Org Lett*. 2003;5(10):1629–32.
46. Anderegg RJ, Betz R, Carr SA, Crabb JW, Duntze W. Structure of *Saccharomyces cerevisiae* mating hormone a-factor. Identification of S-farnesyl cysteine as a structural component. *J Biol Chem*. 1988;263(34):18236–40.
47. Rico J, Pardo E, Orejas M. Enhanced production of a plant monoterpene by overexpression of the 3-hydroxy-3-methylglutaryl coenzyme A reductase catalytic domain in *Saccharomyces cerevisiae*. *Appl Environ Microbiol*. 2010;76(19):6449–54.
48. Szappanos B, Kovács K, Szamecz B, Honti F, Costanzo M, Baryshnikova A, Gelius-Dietrich G, Lercher MJ, Jelasity M, Myers CL, et al. An integrated approach to characterize genetic interaction networks in yeast metabolism. *Nat Genet*. 2011;43(7):656–62.
49. Li BZ, Cheng JS, Ding MZ, Yuan YJ. Transcriptome analysis of differential responses of diploid and haploid yeast to ethanol stress. *J Biotechnol*. 2010;148(4):194–203.
50. Villa-Garcia MJ, Choi MS, Hinz FI, Gaspar ML, Jesch SA, Henry SA. Genome-wide screen for inositol auxotrophy in *Saccharomyces cerevisiae* implicates lipid metabolism in stress response signaling. *Mol Genet Genomics*. 2011;285(2):125–49.
51. Ling H, Chen B, Kang A, Lee JM, Chang MW. Transcriptome response to alkane biofuels in *Saccharomyces cerevisiae*: identification of efflux pumps involved in alkane tolerance. *Biotechnol Biofuels*. 2013;6(1):95.
52. Chumnanpuen P, Nookaew I, Nielsen J. Integrated analysis, transcriptome-lipidome, reveals the effects of INO-level (INO2 and INO4) on lipid metabolism in yeast. *BMC Syst Biol*. 2013;7(Suppl 3):S7.
53. Vemuri GN, Eiteman MA, McEwen JE, Olsson L, Nielsen J. Increasing NADH oxidation reduces overflow metabolism in *Saccharomyces cerevisiae*. *Proc Natl Acad Sci USA*. 2007;104(7):2402–7.
54. Zhang C, Chen X, Zou R, Zhou K, Stephanopoulos G, Too HP. Combining genotype improvement and statistical media optimization for isoprenoid production in *E. coli*. *PLoS ONE*. 2013;8(10):e75164.

Submit your next manuscript to BioMed Central and we will help you at every step:

- We accept pre-submission inquiries
- Our selector tool helps you to find the most relevant journal
- We provide round the clock customer support
- Convenient online submission
- Thorough peer review
- Inclusion in PubMed and all major indexing services
- Maximum visibility for your research

Submit your manuscript at
www.biomedcentral.com/submit

